

Terapevticheskii Arkhiv 2018 vol.90 N8, pages 13-26

Russian consensus on exo- and endocrine pancreatic insufficiency after surgical treatment

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Media Sphera Publishing Group. All rights reserved. The Russian consensus on exo- and endocrine pancreatic insufficiency after surgical treatment was prepared on the initiative of the Russian "Pancreatic Club" on the Delphi method. His goal was to clarify and consolidate the opinions of specialists on the most relevant issues of diagnosis and treatment of exo- and endocrine insufficiency after surgical interventions on the pancreas. An interdisciplinary approach is provided by the participation of leading gastroenterologists and surgeons.

<http://dx.doi.org/10.26442/terarkh201890813-26>

Keywords

Consensus, Endocrine pancreatic insufficiency, Exocrine pancreatic insufficiency, Pancreatic Club, Pancreatic surgery

References

- [1] Andren-Sandberg A, Hardt PD. Second Giessen International Workshop on Interactions of Exocrine and Endocrine Pancreatic Diseases. Castle of Rauischholzhausen of the Justus-Liebig-university, Giessen (Rauischholzhausen), Germany. March 7-8, 2008. JOP. 2008 Jul 10;9(4):541-575.
- [2] Khatkov IE, Maev IV, Abdulkhakov SR, et al. The Russian consensus on the diagnosis and treatment of chronic pancreatitis. Terapevticheskii Arkhiv. 2017;89(2):105-113 (In Russ.) doi: 10.17116/terarkh2017892105-113
- [3] American Diabetes Association. Diagnosis and classification of diabetes mellitus. Diabetes Care. 2014;37 Suppl 1:S81-90. doi: 10.2337/dc14-S081
- [4] Sabater L, Ausania F, Bakker OJ, et al Evidence-based guidelines for the management of exocrine pancreatic insufficiency after pancreatic surgery. Ann Surg. 2016;264(6):949-958. doi: 10.1097/SLA.0000000000001732
- [5] Linstone H, Turoff M. The Delphi method: Techniques and applications. URL: <http://www.is.njit.edu/pubs/delphibook>
- [6] Symersky T, van Hoorn B, Masclee AA The outcome of a long-term follow-up of pancreatic function after recovery from acute pancreatitis. JOP. 2006 Sep 10;7(5):447-453.
- [7] Rana SS, Bhasin DK, Rao C, et al. Comparative evaluation of structural and functional changes in pancreas after endoscopic and surgical management of pancreatic necrosis. Ann Gastroenterol. 2014;27:162-166.
- [8] Gupta R, Wig JD, Bhasin DK, et al. Severe acute pancreatitis: the life after. J Gastrointest Surg. 2009;13(7):1328-1336. doi: 10.1007/s11605-009-0901-z. Epub 2009 May 5.
- [9] Boreham B, Ammori BJ. A prospective evaluation of pancreatic exocrine function in patients with acute pancreatitis: correlation with extent of necrosis and pancreatic endocrine insufficiency. Pancreatol. 2003;3(4):303-308. doi: 10.1159/000071768
- [10] Garip G, Sarandol E, Kaya E. Effects of disease severity and necrosis on pancreatic dysfunction after acute pancreatitis. World J Gastroenterol. 2013 Nov 28;19(44):8065-8070. doi: 10.3748/wjg.v19.i44.8065
- [11] Omelyanovich DA. Otdalennyye rezul'taty lecheniya tyazhelogo ostrogo pankreatita [Longterm results of treatment of severe acute pancreatitis]. Thesis for the degree of candidate of medical sciences. Moscow; 2015 (In Russ.)
- [12] Bornman PC, Botha JF, Ramos JM, et al. Guideline for the diagnosis and treatment of chronic pancreatitis. S Afr Med J. 2010;100(12):845-860.
- [13] Chakraborty PP, Chowdhury S. A look inside the pancreas: The «Endocrine-exocrine Cross-talk». Endocrinol Metab Synd. 2015;4(1):160-164.
- [14] Pezzilli R, Andriulli A, Bassi C, et al. Exocrine Pancreatic Insufficiency collaborative (EPIC) Group. Exocrine pancreatic insufficiency in adults: A shared position statement of the Italian association for the study of the pancreas. World J Gastroenterol. 2013;19(44):7930-7946. doi: 10.3748/wjg.v19.i44.7930
- [15] Czako L, Hegyi P, Rakonczay Z Jr, et al. Interactions between the endocrine and exocrine pancreas and their clinical relevance. Pancreatol. 2009;9(4):351-359. doi: 10.1159/000181169
- [16] De-Madaria E, Abad-Gonzalez A, Aparicio JR, et al. The Spanish Pancreatic Club's recommendations for the diagnosis and treatment of chronic pancreatitis: part 2 (treatment). Pancreatol. 2013;13(1):18-28. doi: 10.1016/j.pan.2012.11.310
- [17] Shevchenko BF, Babiy AM, Makarchuk VA. Morphofunctional state of the pancreas at the stages of its fibrous transformation in chronic pancreatitis. Visnik problem biologii meditsini = Bulletin of Problems of Biology and Medicine. 2014;4(113):305-312 (In Russ.)
- [18] Matsumoto J, Traverso LW. Exocrine function following the Whipple operation as assessed by stool elastase. J Gastrointest Surg. 2006;10(9):1225-1229. doi: 10.1016/j.gassur.2006.08.001
- [19] Roeyen G, Berrevoet F. Pancreatic exocrine insufficiency after pancreaticoduodenectomy is more prevalent with pancreaticogastrostomy than with pancreaticojejunostomy. A retrospective multicenter observational cohort study. HPB (Oxford). 2016;18(12):1017-1022. doi: 10.1016/j.hpb.2016.09.002
- [20] Sikkens EC, Cahen DL, de Wit J, et al. A prospective assessment of the natural course of the exocrine pancreatic function in patients with a pancreatic head tumor. J Clin Gastroenterol. 2014;48(5):e43-e46. doi: 10.1097/MCG.0b013e31829f56e7
- [21] Yuasa Y, Murakami Y, Nakamura H, Uemura K, Ohge H, Sudo T, Hashimoto Y, Nakashima A, Hiyaama E, Sueda T. Histological loss of pancreatic exocrine cells correlates with pancreatic exocrine function after pancreatic surgery. Pancreas. 2012;41(6):928-933. doi: 10.1097/MPA.0b013e31823d837d
- [22] Halloran CM, Cox TF, Chauhan S, et al. Partial pancreatic resection for pancreatic malignancy is associated with sustained pancreatic exocrine failure and reduced quality of life: a prospective study. Pancreatol. 2011;11:535-545. doi: 10.1159/000333308
- [23] Speicher JE, Traverso LW. Pancreatic exocrine function is preserved after distal pancreatectomy. PMID: 20387129. doi: 10.1007/s11605-010-1184-0

- [24] Bartel MJ, Asbun H, Stauffer J, Raimondo M. Pancreatic exocrine insufficiency in pancreatic cancer: A review of the literature. *Dig Liver Dis.* 2015;47(12):1013-1020. doi: 10.1016/j.dld.2015.06.015
- [25] Vincent A, Herman J, Schulick R, et al. Pancreatic cancer. *Lancet.* 2011;378(9791):607-620. doi: 10.1016/S0140-6736(10)62307-0
- [26] Hart PA, Chari ST. Diabetes mellitus and pancreatic cancer: why the association matters? *Pancreas.* 2013;42(8):1207-1209. doi: 10.1097/MPA.0b013e3182a7c963
- [27] Muniraj T, Chari ST. Diabetes and pancreatic cancer. *Minerva Gastroenterol Dietol.* 2012;58(4):331-345.
- [28] Schmied BM, Ulrich AB, Matsuzaki H, et al. Alteration of the Langerhans islets n pancreatic cancer patients. *Int J Pancreatol.* 2000;28:187-197.
- [29] Wu X, Zhang Q, Wang X, et al. Survivin is required for beta-cell mass expansion in the pancreatic duct-ligated mouse model. *PLoS One.* 2012;7(8):e41976. doi: 10.1371/journal.pone.0041976
- [30] Partelli S, Frulloni L, Minniti C, et al. Faecal elastase-1 is an independent predictor of survival in advanced pancreatic cancer. *Dig Liver Dis.* 2012;44(11):945-951. doi: 10.1016/j.dld.2012.05.017. Epub 2012 Jun 28.
- [31] Tseng DS, Molenaar IQ, Besselink MG, et al. Pancreatic Exocrine Insufficiency in Patients With Pancreatic or Periapillary Cancer: A Systematic Review. *Pancreas.* 2016;45(3):325-330. doi: 10.1097/MPA.0000000000000473
- [32] Ghaneh P, Neoptolemos JP. Exocrine pancreatic function following pancreatectomy. *Ann N Y Acad Sci.* 1999;880:308-318.
- [33] Lemaire E, o'Toole D, Sauvanet A, et al. Functional and morphological changes in the pancreatic remnant following pancreaticoduodenectomy with pancreaticogastric anastomosis. *Br J Surg.* 2000;87(4):434-438.
- [34] Dominguez-Munoz JE. Pancreatic exocrine insufficiency: diagnosis and treatment. *J Gastroenterol Hepatol.* 2011;26 Suppl 2:12-16. doi: 10.1111/j.1440-1746.2010.06600.x
- [35] Kahl S, Malfertheiner P. Exocrine and endocrine pancreatic insufficiency after pancreatic surgery. *Best Pract Res Clin Gastroenterol.* 2004;18(5):947-955.
- [36] Yang CJ, Bliss LA, Schapira EF, et al. Systematic review of early surgery for chronic pancreatitis: impact on pain, pancreatic function, and re-intervention. *J Gastrointest Surg.* 2014;18(10):1863-1869. doi: 10.1007/s11605-014-2571-8
- [37] Vinokurova LV, Bordin DS, Dubtsova EA, et al. Some aspects of treatment of patients after operations on the pancreas. *Consilium Medicum.* 2015;8:38-41 (In Russ.)
- [38] Phillips ME. Pancreatic exocrine insufficiency following pancreatic resection. *Pancreatol.* 2015;15(5):449-455. doi: 10.1016/j.pan.2015.06.003
- [39] Morera-ocon FJ, Sabater-orti L, Munoz-Fornier E, et al. Considerations on pancreatic exocrine function after pancreaticoduodenectomy. *World J Gastrointest Oncol.* 2014;6(9):325-329. doi: 10.4251/wjgo.v6.i9.325
- [40] Kang CM, Lee JH. Pathophysiology after pancreaticoduodenectomy. *World J Gastroenterol.* 2015;21(19):5794-5804. doi: 10.3748/wjg.v21.i19.5794
- [41] Lim PW, Dinh KH, Sullivan M, et al. Thirty-day outcomes underestimate endocrine and exocrine insufficiency after pancreatic resection. *HPB.* 2016;18(4):360-366. doi: 10.1016/j.hpb.2015.11.003
- [42] Nakamura H, Murakami Y, Uemura K, et al. Predictive factors for exocrine pancreatic insufficiency after pancreatoduodenectomy with pancreaticogastrostomy. *J Gastrointest Surg.* 2009;13(7):1321-1327. doi: 10.1007/s11605-009-0896-5
- [43] Valiev RK, Fainstein IA, Nechushkin NI, Nurburdyev MB. Evaluation of the exocrine function of the pancreas after pancreatoduodenal resection. *Bulletin of the FGBU RCRC n.a. NN Blokhin.* 2015;26(1):59-61 (In Russ.)
- [44] Lyadov VK, Novozhilov NV. Drug therapy after operations on the pancreas: a review of the literature. *Gastroenterologiya = Gastroenterology.* 2015;(2):21-25 (In Russ.)
- [45] Nakamura H, Murakami Y, Uemura K, et al. Reduced pancreatic parenchymal thickness indicates exocrine pancreatic insufficiency after pancreatoduodenectomy. *J Surg Res.* 2011;171(2):473-478. doi: 10.1016/j.jss.2010.03.052
- [46] Russian Society of Surgeons. Association of hepatopancreatobiliary surgeons of CIS countries. Clinical recommendations for surgical treatment of patients with chronic pancreatitis, 2014 (In Russian) <http://xn-9sbdbejx7bdduahou3a5d.xn-p1ai/stranicaipravlenija/unkr/abdominalnaja-hirurgija/klinicheskie-rekomendacipo-hirurgicheskomu-lecheniyu-bolnyh-hronicheskim-pankreatitom.html>
- [47] Frulloni L, Falconi M, Gabbriellini A, et al. Italian consensus guidelines for chronic pancreatitis. *Dig Liver Dis.* 2010;42 Suppl 6:S381-406. doi: 10.1016/S1590-8658(10)60682-2
- [48] Bachmann K, Tomkoetter L, Erbes J, et al. Beger and Frey procedures for treatment of chronic pancreatitis: comparison of outcomes at 16-year follow-up. *J Am Coll Surg.* 2014;219(2):208-216. doi: 10.1016/j.jamcollsurg.2014.03.040
- [49] Muller MW, Friess H, Martin DJ, et al. Long-term follow-up of a randomized clinical trial comparing Beger with pylorus-preserving Whipple procedure for chronic pancreatitis. *Br J Surg.* 2008;95(3):350-356.
- [50] Strate T, Bachmann K, Busch P, et al. Resection vs drainage in treatment of chronic pancreatitis: long-term results of a randomized trial. *Gastroenterology.* 2008;134(5):1406-1411. doi: 10.1053/j.gastro.2008.02.056
- [51] Zhou Y, Zhao M, Wu L, et al. Short- and long-term outcomes after enucleation of pancreatic tumors: An evidence-based assessment. *Pancreatol.* 2016;16(6):1092-1098. doi: 10.1016/j.pan.2016.07.006
- [52] Jilesen AP, van Eijck CH, Busch oR, et al. Postoperative outcomes of Enucleation and Standard Resections in Patients with a Pancreatic Neuroendocrine Tumor. *World J Surg.* 2016;40(3):715-728. doi: 10.1007/s00268-015-3341-9
- [53] Crippa S, Bassi C, Warshaw AL, et al. Middle pancreatectomy: indications, short- and long-term operative outcomes. *Ann Surg.* 2007;246(1):69-76. doi: 10.1097/01.sla.0000262790.51512.57
- [54] Keim V, Klar E, et al. Postoperative care following pancreatic surgery: surveillance and treatment. *Deut Arzteblatt Intern.* 2009;106(48):789-794. doi: 10.3238/arztebl.2009.0789
- [55] Kang JS, Jang JY, Kang MJ, et al. Endocrine Function Impairment After Distal Pancreatectomy: Incidence and Related Factors. *World J Surg.* 2016;40(2):440-446. doi: 10.1007/s00268-01-3228-9
- [56] Dominguez-Munoz JE, Iglesias-Garcia J, Vilarino-Insua M, Iglesias-Rey M. C-mixed triglyceride breath test to assess oral enzyme substitution therapy in patients with chronic pancreatitis. *Clin Gastroenterol Hepatol.* 2007;5(4):484-488. doi: 10.1016/j.cgh.2007.01.004
- [57] Ivashkin VT, Maev IV, okhlobystin AV, et al. Guidelines of the Russian gastroenterological association on diagnostics and treatment of a chronic pancreatitis. *Rossiyskiy Zhurnal Gastroenterologii, Hepatologii i Koloproktologii = Russian Journal of Gastroenterology, Hepatology, Coloproctology.* 2014; 24(4):70-97 (In Russ.)
- [58] Lohr JM. Exocrine pancreatic insufficiency. *Bremen: UNI-MED;* 2010. 91 p.
- [59] Sikkens EC, Cahen DL, van Eijck C, et al. Patients with exocrine insufficiency due to chronic pancreatitis are undertreated: a Dutch national survey. *Pancreatol.* 2012;12(1):71-73. doi: 10.1016/j.pan.2011.12.010
- [60] Forsmark CE. Chronic pancreatitis and malabsorption. *Am J Gastroenterol.* 2004;99(7):1355-1357. doi: 10.1111/j.1572-0241.2004.70661.x
- [61] Kucheryavyy YuA, Andreev DN. Nutritional status in patients with chronic pancreatitis. *J Nutr Ther.* 2014;3(3):122-132.
- [62] Nils Ewald N, Hardt PD. Diagnosis and treatment of diabetes mellitus in chronic pancreatitis. *World J Gastroenterol.* 2013;19(42):7276-7281. doi: 10.3748/wjg.v19.i42.7276
- [63] Rickels MR, Bellin M, Toledo FG, et al. Pancreas Fest Recommendation Conference Participants. Detection, evaluation and treatment of diabetes mellitus in chronic pancreatitis: recommendations from Pancreas Fest 2012. *Pancreatol.* 2013;13(4):336-342. doi: 10.1016/j.pan.2013.05.002
- [64] Meier JJ, Giese A. Diabetes Associated With Pancreatic Diseases. *Curr opin Gastroenterol.* 2015;31(5):400-406. doi: 10.1097/MoG.0000000000000199
- [65] Makuk J. Management of pancreaticogenic diabetes: challenges and solutions. *Diabetes Metab Syndr Obes.* 2016;9:311-315. doi: 10.2147/DMSO.S99701. eCollection 2016
- [66] Olson DE, Rhee MK, Herrick K, et al. Screening for diabetes and prediabetes with proposed A1C-based diagnostic criteria. *Diabetes Care.* 2010;33(10):2184-2189. doi: 10.2337/dc10-0433
- [67] Kondrup J, Rasmussen HH, Hamborg o, Stanga Z. Nutritional risk screening (NRS 2002): a new method based on an analysis of controlled clinical trials. *Clin Nutr.* 2003;22:321-336.
- [68] Sorensen J, Kondrup J, Prokopowicz J, et al. EurooopS: an international, multicentre study to implement nutritional risk screening and evaluate clinical outcome. *Clin Nutr.* 2008;27(3):340-349. doi: 10.1016/j.clnu.2008.03.012
- [69] ockenga J. Importance of nutritional management in diseases with exocrine pancreatic insufficiency. *HPB (Oxford).* 2009;11 Suppl 3:11-15. doi: 10.1111/j.1477-2574.2009.00134.x
- [70] Goess R, Ceyhan Go, Friess H. Pancreatic exocrine insufficiency after pancreatic surgery. *Panminerva Med.* 2016;58(2):151-159.
- [71] Hashimoto D, Chikamoto A, ohmura M, et al. Impact of postoperative weight loss on survival after resection for pancreatic cancer. *JPN.* 2015;39(5):598-603. doi: 10.1177/0148607114520992
- [72] Augustin T, Burstein MD, Schneider EB, et al. Frailty predicts risk of life-threatening complications and mortality after pancreatic resections. *Surgery.* 2016;160(4):987-996. doi: 10.1016/j.surg.2016.07.010
- [73] onesti JK, Wright GP, Kenning SE, et al. Sarcopenia and survival in patients undergoing pancreatic resection. *Pancreatol.* 2016;16(2):284-289. doi: 10.1016/j.pan.2016.01.009
- [74] Lohr JM, oliver MR, Frulloni L. Synopsis of recent guidelines on pancreatic exocrine insufficiency. *UEGJ.* 2013;1(2):79-83. doi: 10.1177/2050640613476500
- [75] Earthman C, Traughber D, Dobratz J, Howell W. Bioimpedance spectroscopy for clinical assessment of fluid distribution and body cell mass. *Nutr Clin Pract.* 2007;22(4):389-405. doi: 10.1177/0115426507022004389
- [76] Mulasi U, Kuchnia AJ, Cole AJ, Earthman CP. Bioimpedance at the bedside: current applications, limitations, and opportunities. *Nutr Clin Pract.* 2015;30(2):180-193. doi: 10.1177/0884533614568155
- [77] Armstrong T, Strommer L, Ruiz-Jasbon F, et al. Pancreaticoduodenectomy for peri-ampullary neoplasia leads to specific micronutrient deficiencies. *Pancreatol.* 2007;7(1):37-44. doi: 10.1159/000101876
- [78] Duggan SN, Smyth ND, o'Sullivan M, et al. The prevalence of malnutrition and fat-soluble vitamin deficiencies in chronic pancreatitis. *Nutr Clin Prac.* 2014;29(3):348-354. doi: 10.1177/0884533614528361
- [79] Nakamura T, Takebe K, Imamura K, et al. Fatsoluble vitamins in patients with chronic pancreatitis (pancreatic insufficiency). *Acta Gastroenterol Belg.* 1996;59(1):10-14.
- [80] Klapdor S, Richter E, Klapdor R. Vitamin D status and per-oral vitamin D supplementation in patients suffering from chronic pancreatitis and pancreatic cancer disease. *Anticancer Res.* 2012;32(5):1991-1998.
- [81] Lindkvist B, Dominguez-Munoz JE, Luaces-Regueira M, et al. Serum nutritional markers for prediction of pancreatic exocrine sufficiency in chronic pancreatitis. *Pancreatol.* 2012;12(4):305-310. doi: 10.1016/j.pan.2012.04.006
- [82] Park JS, Chung HK, Hwang HK, et al. Postoperative nutritional effects of early enteral feeding compared with total parental nutrition in pancreaticoduodenectomy patients: a prospective, randomized study. *J Korean Med Sci.* 2012;27(3):261-267. doi: 10.3346/jkms.2012.27.3.261

- [83] Berry AJ. Pancreatic surgery: indications, complications, and implications for nutrition intervention. *Nutr Clin Pract.* 2013;28(3):330-357. doi: 10.1177/0884533612470845
- [84] Kucheryavy YuA, Maev IV, Moskaleva AB, et al. Influence of the nutritional status on the course of chronic pancreatitis. *Medical Council.* 2012;(2):100-104 (In Russ.)
- [85] Dujsikova H, Dite P, Tomandl J, et al. occurrence of metabolic osteopathy in patients with chronic pancreatitis. *Pancreatol.* 2008;8(6):583-586. doi: 10.1159/000159845
- [86] Chernyshova IV, Drozdov VN, Vinokurova LV, et al. Exocrine pancreatic insufficiency and bone mineral density disorders in patients with chronic pancreatitis. *Vestnik Kluba Pankreatologov = Bulletin of the Club of Pancreatologists.* 2010;(1):52-54 (In Russ.)
- [87] Duggan SN, o'Sullivan M, Hamilton S, et al. Patients with chronic pancreatitis are at increased risk for osteoporosis. *Pancreas.* 2012;41(7):1119-1124. doi: 10.1097/MPA.0b013e31824abb4d
- [88] Prado CM, Lieffers JR, McCargar LJ, et al. Prevalence and clinical implications of sarcopenic obesity in patients with solid tumours of the respiratory and gastrointestinal tracts. A populationbased study. *Lancet Oncol.* 2008;9(7):629-635. doi: 10.1016/S1470-2045(08)70153-0
- [89] Mourtzakis M, Prado CM, Lieffers JR, et al. A practical and precise approach to quantification of body composition in cancer patients using computed tomography images acquired during routine care. *Appl Physiol Nutr Metab.* 2008;33(5):997-1006. doi: 10.1139/H08-075
- [90] Gomez-Perez SL, Haus JM, Sheean P, et al. Measuring Abdominal Circumference and Skeletal Muscle From a Single Cross-Sectional Computed Tomography Image: A Step-by-Step Guide for Clinicians Using National Institutes of Health Image. *J Parenter Enteral Nutr.* 2016;40(3):308-318. doi: 10.1177/0148607115604149
- [91] Kilgour RD, Cardiff K, Rosenthal L, et al. Use of prediction equations to determine the accuracy of whole-body fat and fat-free mass and appendicular skeletal muscle mass measurements from a single abdominal image using computed tomography in advanced cancer patients. *Appl Physiol Nutr Metab.* 2016;41(1):70-75. doi: 10.1139/apnm-2015-0068
- [92] Baker ST, Strauss BJ, Prendergast LA, et al. Estimating dual-energy X-ray absorptiometry-derived total body skeletal muscle mass using single-slice abdominal magnetic resonance imaging in obese subjects with and without diabetes: a pilot study. *Eur J Clin Nutr.* 2012;66(5):628-632. doi: 10.1038/ejcn.2012.3
- [93] Hoffmeister A, Mayerle J, Beglinger C, et al. English language version of the S3-consensus guidelines on chronic pancreatitis: Definition, aetiology, diagnostic examinations, medical, endoscopic and surgical management of chronic pancreatitis. *Z Gastroenterol.* 2015;53(12): 1447-1495. doi: 10.1055/s-0041-107379
- [94] Smith RC, Smith SF, Wilson J, et al. Summary and recommendations from the Australasian guidelines for the management of pancreatic exocrine insufficiency. Working Party of the Australasian Pancreatic Club. *Pancreatol.* 2016;16(2):164-180. doi: 10.1016/j.pan.2015.12.006
- [95] Khatkov IE, Maev IV, Bordin DS, et al. The Russian consensus on the diagnosis and treatment of chronic pancreatitis: Enzyme replacement therapy. *Terapevticheskiy Arkhiv.* 2017;89(8):80-87 (In Russ.) doi: 10.17116/terarkh201789880-87
- [96] Somaraju UR, Solis-Moya A. Pancreatic enzyme replacement therapy for people with cystic fibrosis. *Cochrane Database Syst Rev.* 2014;10:CD008227. doi: 10.1002/14651858.CD008227.pub2
- [97] Dominguez-Munoz JE. Pancreatic enzyme replacement therapy: exocrine pancreatic insufficiency after gastrointestinal surgery. *HPB (Oxford).* 2009;11 Suppl 3:3-6. doi: 10.1111/j.1477-2574.2009.00132.x
- [98] Conwell DL, Lee LS, Yadav D, et al. American Pancreatic Association Practice Guidelines in Chronic Pancreatitis: evidence-based report on diagnostic guidelines. *Pancreas.* 2014;43(8):1143-1162. doi: 10.1097/MPA.0000000000000237
- [99] Ito T, Ishiguro H, ohara H, et al. Evidence-based clinical practice guidelines for chronic pancreatitis 2015. *J Gastroenterol.* 2016;51(2):85-92. doi: 10.1007/s00535-015-1149-x
- [100] Sadeghi N, Abbruzzese JL, Yeung SC, et al. Metformin use is associated with better survival of diabetic patients with pancreatic cancer. *Clin Cancer Res.* 2012;18:2905-2912.
- [101] Delhaye M, van Steenberghe W, Csemeli E, et al. Belgian consensus on chronic pancreatitis in adults and children: statements on diagnosis and nutritional, medical, and surgical treatment. *Acta Gastroenterol Belg.* 2014;77(1):47-65.
- [102] Forsmark CE. Management of Chronic Pancreatitis. *Gastroenterology.* 2013;144(6):1282-1291.e3. doi: 10.1053/j.gastro.2013.02.008
- [103] Karpoff H, Klimstra DS, Brennan MF, Conlon KC. Results of total pancreatectomy for adenocarcinoma of the pancreas. *Arch Surg.* 2001;136(1):44-47.
- [104] Egorov VI, Vishnevsky VA, Kovalenko ZA, et al. Total pancreatectomy. *Khirurgiya. Zhurnal imeni N.I. Pirogova = Surgery. Journal n.a. N.I. Pirogov.* 2012;(7):85-92 (In Russ.)
- [105] Kucheryavy YuA. External secretory pancreatic insufficiency after surgical interventions on the stomach and pancreas. *Khirurgiya. Prilozhenie k zhurnalu Consilium Medicum = Surgery. Supplement to the journal Consilium Medicum.* 2012;(1):42-46 (In Russ.)
- [106] Tran TC, van Hof G, Kazemier G, et al. Pancreatic fibrosis correlates with exocrine pancreatic insufficiency after pancreatoduodenectomy. *Dig Surg.* 2008;25(4):311-318. doi: 10.1159/000158596
- [107] Ito K. Duodenum preservation in pancreatic head resection to maintain pancreatic exocrine function (determined by pancreatic function diagnostant test and cholecystokinin secretion). *J Hepatobiliary Pancreat Surg.* 2005;12(2):123-128. doi: 10.1007/s00534-004-0954-z